

2+1 Gravity without Dynamics

Viqar Husain

International Centre for Theoretical Physics

PO Box 586, 34100 Trieste, Italy*

and

Department of Physics

University of Utah, Salt Lake City, UT 84112, USA

Abstract

A three dimensional generally covariant theory is described that has a 2+1 canonical decomposition in which the Hamiltonian constraint, which generates the dynamics, is absent. Physical observables for the theory are described and the classical and quantum theories are compared with ordinary 2+1 gravity.

* present address. email: viqar@itsictp.bitnet

The work on 3+1 gravity in recent years using the Ashtekar canonical variables [1-3] has prompted the study of a number of toy models that have Hamiltonian formulations similar to those for general relativity. Among these models are 2+1 gravity[4], and a 4D generally covariant theory that has the peculiar property that its canonical decomposition does not contain the Hamiltonian constraint[5]. This latter theory is constructed from an $SU(2)$ connection a , and a dreibein e with internal indices in $SU(2)$ (rather than the usual $SL(2,C)$), and its action is $\int_M Tr(e \wedge e \wedge F[a])$. In the canonical decomposition of this theory, the phase space and the other constraints, the 3D spatial diffeomorphism and Gauss law constraints, have exactly the same form as those for canonical general relativity written in terms of the Ashtekar variables. The other difference between this 4D model and general relativity is that the phase space coordinate is a real $SU(2)$ connection for the former and a complex one for the latter. Therefore the analog of the reality conditions present in the Ashtekar formulation are easy to implement in the quantum theory.

The reason that the Hamiltonian constraint (and hence ‘dynamics’) is absent in this model is essentially that the action is constructed from a dreibein, so that the spacetime metric is degenerate with signature $(0, +, +, +)$. It turns out (via the equations of motion) that the degeneracy direction is also a Killing direction. Thus the 4D theory in its generally covariant form is effectively already ‘dimensionally reduced’.

In this letter, a 3D generally covariant theory is described that has the same relationship to 3D gravity that the above mentioned model has to 4D gravity. The theory is topological, as is 3D gravity, and is ‘exactly soluble’ in the same sense, but with the dynamics really absent. (One might be tempted to say that there is no dynamics in 3D gravity as well. This is not the case however, since in the standard initial value formulation, the geometric variables (the spatial metric and the extrinsic curvature) do evolve via the hamiltonian constraint). A comparison of the classical and quantum theories with those of 3D gravity is therefore useful, and may help to clarify conceptual issues regarding dynamical versus non-dynamical observables that have been discussed recently in the

literature[6-8]. It may also help in identifying a natural intrinsic time variable for 2+1 gravity.

In order that there be no Hamiltonian constraint (and hence no ‘dynamics’) in the Hamiltonian theory, we follow the example of the 4D theory[5] and seek a generally covariant 3D action that is constructed from a zweibein and a gauge field. Since Wit ISO(2,1) Chern-Simons theory [9], (which has a dreibein built into it), a natural choice for an action with a zweibein is provided by the an ISO(2) Chern-Simons theory. ISO(2) gives a degenerate spacetime metric with signature $(0, +, +)$ as desired, (although the latter may be of interest as well). There is however a problem with this since there is no non-degenerate quadratic form on these Lie algebras. This is rectified by considering the Lie algebras with a cosmological constant λ . The generators P_i and J , with non-zero λ now satisfy the commutation rules

$$[P_i, P_j] = \lambda \epsilon_{ij} J \quad [J, P_i] = \epsilon_{ij} P^j \quad (1)$$

where the (internal) indices $i, j = 1, 2$ are raised by the 2D metrics $(+, +)$ for λ positive and $(-, +)$ for λ negative. The algebras for $\lambda = +-$ are $\text{so}(3)$ and $\text{so}(2,1)$ respectively, corresponding to the DeSitter groups of Euclidean or Minkowskian 2D space. The relevant connection is $A_\alpha \equiv e_\alpha^i P^i + \omega_\alpha J$ where $\alpha, \beta \dots$ are the spacetime indices. The Chern-Simons action

$$I = \int Tr(A \wedge dA + \frac{2}{3} A \wedge A \wedge A) \quad (2)$$

with the trace $\langle J, J \rangle = 1$, $\langle P_i, P_j \rangle = \lambda \delta_{ij}$ and $\langle J, P_i \rangle = 0$ reduces to

$$I = \int_M \epsilon^{\alpha\beta\gamma} [\omega_\alpha \partial_\beta \omega_\gamma + \lambda e_\alpha^i \partial_\beta e_\gamma^i + \lambda \epsilon_{ij} e_\alpha^i e_\beta^j \omega_\gamma] \quad (3)$$

The degenerate spacetime metric is $g_{\alpha\beta} = e_\alpha^i e_\beta^i$ and the degeneracy direction is given by the vector density $n^\alpha = \epsilon^{\alpha\beta\gamma} e_\beta^i e_\gamma^j \epsilon_{ij}$. We have the relation $g_{\alpha\beta} n^\alpha = 0$. A straightforward 2+1 decomposition, assuming the spacetime topology $\Sigma \times R$, yields the canonically conjugate phase space variables (e_1^i, e_2^i) and (ω_1, ω_2) satisfying the commutation rules

$\{e_a^i(x), e_b^j(y)\} = (1/\lambda)\epsilon_{ab}\delta^{ij}\delta^2(x-y)$ and $\{\omega_a(x), \omega_b(y)\} = \epsilon_{ab}\delta^2(x-y)$. The indices $a, b, ..$ are the spatial projections of the spacetime indices. There are three constraints found by varying the action with respect to the time components, ω_0 and e_0^i :

$$g \equiv \epsilon^{ab}(\partial_a \omega_b + \frac{\lambda}{2}\epsilon_{ij}e_a^i e_b^j) = 0 \quad (4)$$

and

$$V^i \equiv \epsilon^{ab}(\partial_a e_b^i + \epsilon_j^i e_a^j \omega_b) = 0 \quad (5)$$

These satisfy the Poisson bracket relations

$$\{g(\Lambda), g(\Lambda')\} = 0 \quad \{g(\Lambda), V(m^i)\} = V(\Lambda \epsilon^{ij} m^j) \quad (6)$$

and

$$\{V(m^i), V(n^i)\} = g(\epsilon^{ij} m^i n^j) \quad (7)$$

where $g(\Lambda) = \int_{\Sigma} \Lambda g$ and $V(m^i) = \int_{\Sigma} m^i V^i$. These are all the (first class) constraints and there are no other constraints.

The first (4) generates internal gauge transformations: rotations of the zweibein $\{g(\Lambda), e_a^i\} = \Lambda \epsilon^{ji} e_a^j$, and Abelian gauge transformations on ω_a , $\{g(\Lambda), \omega_a\} = \partial_a \Lambda$. The second (5) is replaced by the linear combination $C_a \equiv \lambda e_a^i V^i + \omega_a g$. It is easy to verify, for example, that $\{\int_{\Sigma} N^a C_a, e_a^i\} = \mathcal{L}_N e_a^i$, where $\mathcal{L}_N$ denotes the Lie derivative with respect to the vector field N^a . It is clear that there is no analog of a Hamiltonian constraint (as desired). By contrast, for 2+1 gravity in the dreibein-spin connection variables, there is an SO(2,1) Gauss law constraint and a vanishing curvature condition for the SO(2,1) connection [9,4]. The latter are three conditions which do contain the Hamiltonian constraint (as well as the two spatial diffeomorphisms C_a).

The gauge invariant physical observables can be readily constructed by noting that the constraints g, V^i are just pieces of the vanishing curvature condition $F_{ab} = V^i P^i + gJ = 0$ for the connection $A_a = e_a^i P^i + \omega_a J$. The observables are therefore just the Wilson lines for

this A , $T[\gamma] \equiv \text{Tr} [\text{Pexp} \int_{\gamma} dx A]$, where the trace is taken over a two by two representation of the generators.

We now restrict attention to the case of $\lambda = 1$ where the algebra (1) is $\text{so}(3)$, since this is the case relevant to spatial metrics of signature $(+, +)$ (and hence to comparisons with 2+1 gravity). The case of $\lambda = -1$ may be similarly treated. Using the fundamental Poisson brackets and the trace identity $\sum_i \text{Tr}[A\tau^i]\text{Tr}[B\tau^i] = \text{Tr}[AB] - \text{Tr}[AB^{-1}]$ (where A, B are $\text{SO}(3)$ matrices and τ^i are the generators), the Wilson line observables satisfy the closed Poisson bracket relations

$$\{T[\alpha], T[\beta]\} = \Delta(\alpha, \beta)(T[\alpha\circ\beta] - T[\alpha\circ\beta^{-1}]) \quad (8)$$

where $\alpha\circ\beta$ denotes composition of the loops α and β and $\Delta(\alpha, \beta)$ is $\pm$ depending on the right/left handed sense of the tangent vectors to the loops at their intersection(s).

On the classical reduced phase space, the $T[\alpha]$ are functions of the homotopy class of the loop α . Therefore, for the torus, for example, the reduced phase space is two-dimensional and is coordinatized by $T[a]$ and $T[b]$ where a, b are the (Abelian) generators of the homotopy group of the torus. In contrast, the reduced phase space for 2+1 gravity for the torus is four dimensional. The reason is as follows: whereas one may use Wilson line observables for 2+1 gravity as well [9,10], the connection in this case is $\text{iso}(2,1)$ valued. Therefore, since there are two Casimir invariants for this Lie algebra, each Wilson line results in two independent observables, the mass and the angular momentum. (These two observables per line may be separated naturally into a Wilson line observable for an $\text{so}(2,1)$ connection, and an observable that is linear in the momentum conjugate to this connection [4]). But for the degenerate theory here, the Lie algebra is $\text{so}(3)$ and there is only one Casimir invariant. Therefore there is only one observable per Wilson line.

It is straightforward to construct a representation of the Poisson algebra (8) on functions of loops and thereby obtain the quantum mechanics of the model. Given functions

of loops β , $a[\beta]$, satisfying the condition $a[\beta] = a[\beta^{-1}]$, we define the operator

$$\hat{T}[\alpha]a[\beta] = i\hbar\Delta(\alpha, \beta)(a[\alpha\circ\beta] - a[\alpha\circ\beta^{-1}]). \quad (9)$$

This definition gives a commutator algebra for the $\hat{T}$ that reduces to (8) in the classical limit.

Turning now to the quantum theory for the torus, we consider, for windings n_1, n_2 of the generators, representation space kets $|n_1, n_2\rangle$ ($= |-n_1, -n_2\rangle$). The action (9) of the operators $\hat{T}[\alpha]$ now specializes to

$$\begin{aligned} \hat{T}[a]|n_1, n_2\rangle &= i\hbar n_2(|n_1 + 1, n_2\rangle - |n_1 - 1, n_2\rangle) \\ \hat{T}[b]|n_1, n_2\rangle &= -i\hbar n_1(|n_1, n_2 + 1\rangle - |n_1, n_2 - 1\rangle) \end{aligned} \quad (10)$$

The inner product is $\langle m_1, m_2 | n_1, n_2 \rangle = \delta_{m_1, n_1} \delta_{m_2, n_2}$. This completes the description of the quantum theory.

The two observables discussed above for the torus are truly non-dynamical since there is no Hamiltonian constraint. Furthermore, another reason we expect there to be two is that, for 2+1 gravity, for the torus, the four dimensional phase space can in principle be coordinatized by conjugate Hamiltonian and intrinsic time functions, together with two observables that evolve with respect to this time variable via the conjugate Hamiltonian. The present theory naturally identifies the two observables that are ‘waiting to evolve’ in this sense, via that Hamiltonian and time variable that will convert this theory to 2+1 gravity (after the appropriate extension of the phase space). Or, to put it another way, if one can isolate these two observables on the phase space of 2+1 gravity (for the torus), the two remaining conjugate variables will provide a natural intrinsic time variable and Hamiltonian. A hint as to what observables of 2+1 gravity may correspond to the ones presented here comes from the Poisson algebra (8). Two (of the four) observables for 2+1 gravity have exactly this algebra [4]! H and T , of the four observables, such that the

Poisson brackets $\{H(x), T(y)\} = \delta^2(x - y)$ are satisfied. In principle this procedure should be possible for spacelike surfaces of arbitrary topology.

Another possible direction that may be pursued using this model concerns 2D gravity. The model is in fact one for topological 2D gravity since there is a 2D metric and a 2D diffeomorphism constraint in the canonical analysis. Thus $SO(3)$ Chern-Simons theory may be interpreted in this way as 2D Euclidean (topological) gravity (or 2D Minkowskian gravity for $SO(2,1)$). It would be of interest to see if matter fields can be coupled to it in such a way that it remains non-dynamical in a 3D sense but becomes a 2D generally covariant field theory with local degrees of freedom. This would provide a way of constructing 2D field theories via canonical decompositions of 3D theories that have degenerate spacetime metrics.

I wish to thank Karel Kuchar for helpful discussions, Lee Smolin for comments on the manuscript, and Prof. Abdus Salam, IAEA, and UNESCO for hospitality at the International Centre for Theoretical Physics. This work was supported in part by NSF grant PHY-8907937 to the University of Utah.

References

- [1] A. Ashtekar, Phys. Rev. Lett. 57 (1986) 2244; Phys. Rev. D36 (1987) 1587; New Perspectives in Canonical Gravity (Naples: Bibliopolis, 1988).
- [2] A. Ashtekar and R. S. Tate, Non-perturbative canonical gravity (Lecture Notes) (Inter-University Centre for Astronomy and Astrophysics, Poona, India).
- [3] C. Rovelli, Class. and Quantum Grav. 8 (1991) 1613.
- [4] A. Ashtekar, V. Husain, C. Rovelli, J. Samuel, and L. Smolin, Class. and Quantum Grav. L185 (1989).
- [5] V. Husain and K. Kuchar, Phys. Rev. D42 (1990) 4070.
- [6] C. Rovelli, Phys. Rev. D42 (1990) 2638.

- [7] S. Carlip, Phys. Rev. D42 (1990) 2647.
- [8] V. Moncrief, J. Math. Phys. 31 (1990) 2978.
- [9] E. Witten, Nucl. Phys. B311 (1988) 46.
- [10] S. Martin, Nucl. Phys. B327 (1989)178.